

A theorem on spherically complete valued abelian groups

Franz-Viktor Kuhlmann

12. 6. 1997

Abstract. We give a criterion for a group homomorphism on a valued abelian group to be surjective and to preserve spherical completeness. We apply this to give a criterion for the existence of integration on a valued differential field. Further, we give a criterion for a sum of spherically complete subgroups of a valued abelian group to be spherically complete. This in turn can be used to determine elementary properties of power series fields in positive characteristic.

Let (A, v) be a valued abelian group. That is, $v : A \ni a \mapsto va$ is a map from A onto a totally ordered set with largest element ∞ , satisfying that $va = \infty \Leftrightarrow a = 0$ and $v(a - a') \geq \min\{va, va'\}$. The latter is the **ultrametric triangle law**. We set $v(A) := \{va \mid a \in A\}$.

We recall some definitions. For $a \in A$ and $\alpha \in v(A)$, we define the **ball**

$$B_\alpha(a) := \{a' \in A \mid v(a - a') \geq \alpha\}.$$

It follows from the ultrametric triangle law that

$$B_\alpha(a) \subseteq B_{\alpha'}(a') \quad \text{if and only if} \quad a \in B_{\alpha'}(a') \text{ and } \alpha \geq \alpha'. \quad (1)$$

A set of balls in (A, v) is called a **nest of balls** if it is totally ordered by inclusion. By (1), this is the case as soon as every two balls in the set have a nonempty intersection. The **intersection** of the nest is defined to be the intersection of all of its balls. Further, (A, v) is called **spherically complete** if every nest of balls has a nonempty intersection.

It is well-known that the valued additive group of every power series field and, more generally, of every maximally valued field is spherically complete (cf. [P1], [P2], [PR], and [K1]). For maximally valued fields, this was essentially proved by Kaplansky in [KA], using the notion of “pseudo Cauchy sequence” instead of “nest of balls”.

Theorem: *Let (A, v) and (B, w) be valued abelian groups and $f : A \rightarrow B$ a group homomorphism. Assume that there is a subset $S \subseteq A$ such that:*

a) the assignment

$$vs \mapsto wf(s) \quad (s \in S)$$

defines an order preserving map $\varphi : v(S) \rightarrow w(B)$,

b) for all $a \in A$ and $s \in S$,

$$va \geq vs \Rightarrow wf(a) \geq wf(s),$$

c) for every $b \in B \setminus \{0\}$ there is some $s \in S$ such that

$$w(b - f(s)) > wb.$$

Suppose further that (A, v) is spherically complete. Then f is surjective and (B, w) is spherically complete.

Proof: We will write fa instead of $f(a)$. First, we observe that $w(B) \setminus \{\infty\} \subseteq \varphi(v(S))$: by assumption c), for every $b \in B$, $b \neq 0$, there is some $s \in S$ such that $w(b - fs) > wb$, which implies that $wb = wfs = \varphi vs \in \varphi(v(S))$. Since φ is order preserving by assumption, it is also injective. Therefore, it admits an inverse $\varphi^{-1} : w(B) \setminus \{\infty\} \rightarrow v(S)$.

Take an arbitrary $a \in A$ and some $\alpha \in v(S)$; we claim that

$$f(B_\alpha(a)) \subseteq B_{\varphi\alpha}(fa).$$

Indeed, if $a' \in B_\alpha(a)$ then $v(a - a') \geq \alpha$, and thus $w(fa - fa') = wf(a - a') \geq \varphi\alpha$ by assumption b), showing that $fa' \in B_{\varphi\alpha}(fa)$.

Next, using that (A, v) is spherically complete we wish to show that

$$\forall a \in A \forall \alpha \in v(S) : f(B_\alpha(a)) = B_{\varphi\alpha}(fa) \quad (2)$$

where the latter is a ball in B . Take any $a \in A$, $\alpha \in v(S)$ and $b \in B_{\varphi\alpha}(fa)$. We have to show that $b \in f(B_\alpha(a))$. Let $B_{\alpha_i}(a_i)$, $i \in I$, be a maximal nest of balls in $B_\alpha(a)$ such that $\alpha_i \in v(S)$ and $b \in B_{\varphi\alpha_i}(fa_i)$ for all i . Since the nest $\{B_\alpha(a)\}$ satisfies these conditions, a maximal nest exists by Zorn's Lemma.

Since (A, v) is spherically complete by assumption, there is some $\tilde{a} \in \bigcap_{i \in I} B_{\alpha_i}(a_i)$. Then $f\tilde{a} \in f(B_{\alpha_i}(a_i)) \subseteq B_{\varphi\alpha_i}(fa_i)$ and thus $w(b - f\tilde{a}) \geq \varphi\alpha_i$ for all i . Suppose that $b \neq f\tilde{a}$. Then by assumption c), there is some $s \in S$ such that $w(b - f\tilde{a}) < w((b - f\tilde{a}) - fs) = w(b - f(\tilde{a} + s)) =: \beta'$. We set $a' := \tilde{a} + s$ and observe that $vs = \varphi^{-1}wfs = \varphi^{-1}w(b - f\tilde{a}) \geq \alpha_i$ for all i . Hence, $a' = \tilde{a} + s \in B_{\alpha_i}(\tilde{a}) = B_{\alpha_i}(a_i)$ for all i .

If $b = fa'$ then $b \in f(B_\alpha(a))$ and we are done. Otherwise $\beta' \neq \infty$, and we can set $\alpha' := \varphi^{-1}\beta'$. As $\beta' > w(b - f\tilde{a}) \geq \varphi\alpha_i$, we have that $\alpha' > \alpha_i$ for all i . This shows that $B_{\alpha'}(a')$ is properly contained in all $B_{\alpha_i}(a_i)$. On the other hand, $b \in B_{\beta'}(fa') = B_{\varphi\alpha'}(fa')$. Therefore, we can enlarge our nest of balls by adding $B_{\alpha'}(a')$. As this contradicts the maximality, we conclude that $b = f\tilde{a} \in f(B_\alpha(a))$, as desired.

To show the surjectivity of f , take any $b \in B$. If $b = 0$ then $b = f0 \in f(A)$. So assume that $b \neq 0$. Then by assumption c), we can choose some $a \in S$ such that $\beta := w(b - fa) > wb$. If $fa = b$ then we are done. Otherwise $\beta \neq \infty$ and we can set $\alpha := \varphi^{-1}\beta \in v(S)$, obtaining that $b \in B_\beta(fa) = f(B_\alpha(a))$. This shows the surjectivity. Let us note that $\beta > wb = wfa$ and hence, $\alpha = \varphi^{-1}\beta > \varphi^{-1}wfa = va$. So we have proved, for all $b \in B$:

$$b \in f(S \cup \{0\}) \quad \vee \quad \exists a \in S \exists \alpha \in v(S) : \alpha > va \wedge b \in f(B_\alpha(a)). \quad (3)$$

Finally, let us prove that (B, w) is spherically complete. Take any nest of balls $\{B_{\beta_j}(b_j) \mid j \in J\}$ in (B, w) . We have to show that this nest has a nonempty intersection. We may assume that $\beta_j \neq \infty$ for all $j \in J$; indeed, if $\beta_{j_0} = \infty$ for some $j_0 \in J$, then the intersection is $\{b_{j_0}\} \neq \emptyset$ and there is nothing more to show.

We claim that there exists a nest of balls $\{B_{\alpha_i}(a_i) \mid i \in I\}$ in (A, v) maximal with the property that for every $i \in I$,

$$\alpha_i \in v(S) \quad \text{and there is some } j \in J \text{ such that } B_{\beta_j}(b_j) \subseteq f(B_{\alpha_i}(a_i)).$$

Indeed, we can choose an arbitrary $j \in J$, set $\alpha := \varphi^{-1}\beta_j$ and then, by the surjectivity of f , choose some $a \in A$ such that $fa = b_j$. Then by (2), $B_{\beta_j}(b_j) = f(B_\alpha(a))$. So $\{B_\alpha(a)\}$ is a nest with the required property. Hence, a maximal nest $\{B_{\alpha_i}(a_i) \mid i \in I\}$ with this property exists by Zorn's Lemma. It follows that

$$\bigcap_{j \in J} B_{\beta_j}(b_j) \subseteq \bigcap_{i \in I} f(B_{\alpha_i}(a_i)) = \bigcap_{i \in I} B_{\varphi\alpha_i}(fa_i). \quad (4)$$

We wish to show that

$$f\left(\bigcap_{i \in I} B_{\alpha_i}(a_i)\right) \subseteq \bigcap_{j \in J} B_{\beta_j}(b_j). \quad (5)$$

Suppose the contrary. Since $f(\bigcap_{i \in I} B_{\alpha_i}(a_i)) \subseteq \bigcap_{i \in I} f(B_{\alpha_i}(a_i))$, we must then have that inequality holds in (4) and that there is an element $\tilde{a} \in \bigcap_{i \in I} B_{\alpha_i}(a_i)$ and some $j_0 \in J$ such that $f\tilde{a} \notin B_{\beta_{j_0}}(b_{j_0}) \subsetneq \bigcap_{i \in I} B_{\varphi\alpha_i}(fa_i)$. The inequality implies that $\beta_{j_0} > \varphi\alpha_i$ for all i . As $b_{j_0}, f\tilde{a} \in B_{\varphi\alpha_i}(fa_i)$, we have that $w(b_{j_0} - f\tilde{a}) \geq \varphi\alpha_i$ for all i . By assumption c), there is some $s \in S$ such that $w(b_{j_0} - f\tilde{a}) < w((b_{j_0} - f\tilde{a}) - fs) = w(b_{j_0} - f(\tilde{a} + s)) =: \tilde{\beta}$. As before, we set $a' := \tilde{a} + s$ and observe that $a' \in B_{\alpha_i}(a_i)$ for all i . We set $\beta' := \min\{\tilde{\beta}, \beta_{j_0}\} < \infty$ and $\alpha' := \varphi^{-1}\beta'$. Then $\beta' > \varphi\alpha_i$ and $\alpha' > \alpha_i$ for all i . It follows that $B_{\alpha'}(a')$ is properly contained in all $B_{\alpha_i}(a_i)$ for all i . On the other hand, $\beta_{j_0} \geq \beta'$ and $w(b_{j_0} - fa') = \tilde{\beta} \geq \beta'$ imply that $B_{\beta_{j_0}}(b_{j_0}) \subseteq B_{\beta'}(fa') = f(B_{\alpha'}(a'))$. Therefore, we can enlarge our nest of balls by adding $B_{\alpha'}(a')$. As this contradicts the maximality, we conclude that (5) must hold.

Since (A, v) is spherically complete by assumption, $\bigcap_{i \in I} B_{\alpha_i}(a_i)$ is nonempty. Hence by (5), $\bigcap_{j \in J} B_{\beta_j}(b_j)$ is nonempty, as desired. $\square$

There is also a version of this theorem for ultrametric spaces, where addition is not available ([K3]).

Remarks: Assume that the conditions of the theorem hold. Then in particular, (3) holds for every $b \in B$.

Take any $a \in S$ and $\alpha \in v(S)$ such that $va < \alpha$. It follows that every $a' \in B_\alpha(a)$ has value $va' = \min\{va, \alpha\} = va$. By (2), $fa' \in f(B_\alpha(a)) = B_{\varphi\alpha}(fa)$. Since $\varphi\alpha > wfa$ and $a \in S$, we have that $wfa' = \min\{wfa, w(fa' - fa)\} = wfa = \varphi va = \varphi va'$. Hence, the subset $S \cup B_\alpha(a) \subseteq A$ still has property a). Trivially, it also has properties b) and c). Therefore, every maximal subset $S \subseteq A$ having properties a), b) and c) will contain every ball $B_\alpha(a)$ of the above form. By (3), it will therefore satisfy that

$$f(S \cup \{0\}) = B.$$

On the other hand, if S can be chosen such that $S - S \subseteq S$ (which in fact means that S is a subgroup), then $f : S \rightarrow B$ will be injective. Indeed, if $s, s' \in S$ such that $fs = fs'$, then $f(s - s') = 0$ and $wf(s - s') = \infty$. With $s - s' \in S$ and $0 \in S$, this implies that $v(s - s') = \infty$, i.e., $s = s'$. If in addition for all $a \in A$ there is some $s \in S$ such that $fs = fa$, then $f : S \rightarrow B$ will be an isomorphism of valued groups. This is the situation of application (A) below.

Note that if S contains an element s_0 such that $fs_0 = 0$ (e.g., if $0 \in S$), then by the first part of our proof, $\varphi : v(S) \rightarrow w(B)$ will be an order preserving bijection. For this, it is not necessary that (A, v) be spherically complete.

Applications:

(A) Let (K, D) be a differential field with field of constants $C = \{a \in K \mid Da = 0\}$. Following M. Rosenlicht [R1], a valuation v of K is called a **differential valuation** if it is trivial on C and satisfies

$$\forall a, b \in K : va \geq 0 \wedge vb > 0 \wedge b \neq 0 \Rightarrow v\left(\frac{bDa}{Db}\right) > 0. \quad (6)$$

These conditions imply that

$$\forall a, b \in K \setminus \{0\}, va \neq 0, vb \neq 0 : va \leq vb \Leftrightarrow vDa \leq vDb. \quad (7)$$

If C is a set of representatives for the v -residues of K , then also the converse holds, i.e., v is differential if and only if it satisfies (7), cf. [R1].

We say that (K, D) **admits integration** if D is surjective. If v is a differential valuation on K , then following Rosenlicht [R2], we say that (K, D) **admits asymptotic integration** if for every $b \in K, b \neq 0$, there is some $a \in K$ such that $v(b - Da) > vb$.

Let k be any field and G any ordered abelian group. The **(generalized) power series field** $k((G))$ consists of all formal sums $a = \sum_{g \in G} c_g t^g$ with $c_g \in k$ and well-ordered **support** $\text{supp}(a) = \{g \in G \mid c_g \neq 0\}$. The **canonical valuation** on $k((G))$ is given by $va := \min \text{supp}(a) \in G$ and $v0 := \infty$.

The following is applied in [KK] to prove integration for exponential-logarithmic power series:

Let $k((G))$ be a power series field, endowed with a derivation D for which k is the field of constants. Suppose that the canonical valuation v of K is a differential valuation (with respect to D). Then $(k((G)), D)$ admits integration if and only if it admits asymptotic integration.

Proof: As we have already mentioned, the additive group of the power series field $k((G))$, endowed with the canonical valuation v , is spherically complete. We let S be the subgroup of all power series a in $k((G))$ without constant term (that is, if $a = \sum_{g \in G} c_g t^g$ then $c_0 = 0$). Then $k((G)) = S \oplus k$ and $v(S) = G \setminus \{0\}$. Therefore, (7) implies assumption a) of the theorem.

If $a = s' + c \in k((G))$ with $s' \in S$ and $c \in k$, then $va = vs' < 0$ or $vs' \geq va \geq 0$ (with $vs' = va > 0$ if $c = 0$). In all cases, $vs' \geq va$. Hence if $s \in S$ such that $va \geq vs$, then $vs' \geq vs$ and by (7), $vDa = vDs' \geq vDs$. This shows that also assumption b) holds.

Since for every $a \in k((G))$ there is some $s \in S$ such that $Da = Ds$, assumption c) holds if and only if $(k((G)), D)$ admits asymptotic integration. Now the theorem proves one direction, and the other one is clear since integration implies asymptotic integration. $\square$

In this application, S is a subgroup of the additive group of $k((G))$, having the property that for every $a \in k((G))$ there is some $s \in S$ such that $Ds = Da$. Hence by the remark above, $D : S \rightarrow k((G))$ is an isomorphism of valued abelian groups. In particular, this shows that for every $b \in k((G))$ there is a *unique* $s \in S$ such that $Ds = b$.

On the other hand, S is not a maximal subset of $k((G))$ having properties a), b) and c). Indeed, as $v(S) = G \setminus \{0\}$, we can pick $a \in S$ and $\alpha \in v(S)$ such that $va < \alpha < 0$. Then $a + c \in B_\alpha(a)$ for every $c \in k$ and thus, $B_\alpha(a) \not\subseteq S$. However, S is a maximal *subgroup* having properties a), b) and c).

(B) Let $(\mathcal{A}, v)$ be a valued abelian group and $A_1, \dots, A_n$ be subgroups of A . We call the sum $A_1 + \dots + A_n \subseteq \mathcal{A}$ **pseudo-direct** if for every $a \in A_1 + \dots + A_n$, $a \neq 0$, there are $a_i \in A_i$ such that

$$v \sum_{i=1}^n a_i = \min_{1 \leq i \leq n} va_i \quad \text{and} \quad v \left(a - \sum_{i=1}^n a_i \right) > va.$$

The following is applied in [K1], [K2] to determine elementary properties of the power series field $\mathbb{F}_p((t))$:

Assume that the subgroups (A_i, v) of $(\mathcal{A}, v)$, $1 \leq i \leq n$, are spherically complete. If the sum $A_1 + \dots + A_n \subseteq \mathcal{A}$ is pseudo-direct, then it is also spherically complete.

Proof: We endow the direct product $A := A_1 \times \dots \times A_n$ with the **minimum valuation** by setting

$$v(a_1, \dots, a_n) := \min_{1 \leq i \leq n} va_i.$$

Further, we set $B := A_1 + \dots + A_n \subseteq \mathcal{A}$ and take $f : A \rightarrow B$ to be the group homomorphism defined by $f(a_1, \dots, a_n) := \sum_{i=1}^n a_i$. We let $S \subseteq A$ be the set of all n -tuples $(a_1, \dots, a_n)$ which satisfy that $v \sum_{i=1}^n a_i = \min_{1 \leq i \leq n} va_i$. We note that $v(S) = v(A) \subseteq v(\mathcal{A})$ and that the map φ defined as in assumption a) of the theorem is simply the identity on $v(S)$. Assumption b) of the theorem holds since by the ultrametric triangle law, we always have that $v \sum_{i=1}^n a_i \geq \min_{1 \leq i \leq n} va_i$. Finally, the sum $A_1 + \dots + A_n \subseteq \mathcal{A}$ is pseudo-direct if and only if assumption c) of the theorem holds. As it is well-known that the direct product of finitely many spherically complete valued abelian groups, endowed with the minimum valuation, is again spherically complete (cf. [K1]), our assertion now follows from the theorem. $\square$

In this application, S will in general not be a subgroup of A , and f will in general not be injective. Indeed, the groups A_i may have non-trivial intersection, so two elements of S may be equal up to a permutation of the coördinates a_i .

(C) Let (A, v) be a spherically complete valued abelian group, and $\alpha \in v(A)$. We leave it to the reader to define a canonical valuation on the quotient group $A/B_\alpha(0)$ and to apply the theorem in order to show that with this valuation, $A/B_\alpha(0)$ is spherically complete.

References

- [KA] Kaplansky, I.: *Maximal fields with valuations I*, Duke Math. J. **9** (1942), 303–321
- [K1] Kuhlmann, F.-V.: *Valuation theory of fields, abelian groups and modules*, preprint, Heidelberg (1996), to appear in the “Algebra, Logic and Applications” series (Gordon and Breach), eds. A. Macintyre and R. Göbel
- [K2] Kuhlmann, F.-V.: *Elementary properties of power series fields over finite fields*, conference hand-out, Toronto (January 1997); prepublication in: Structures Algébriques Ordonnées, Séminaire Paris VII (1997), and in: The Fields Institute Preprint Series (June 1997)
- [K3] Kuhlmann, F.-V.: *Ultrametric spaces*, in preparation
- [KK] Kuhlmann, F.-V. – Kuhlmann, S.: *Exponential-logarithmic power series*, in preparation
- [P1] Priß-Crampe, S.: *Angeordnete Strukturen. Gruppen, Körper, projektive Ebenen*, Ergebnisse der Mathematik und ihrer Grenzgebiete **98**, Springer (1983)
- [P2] Priß-Crampe, S.: *Der Banachsche Fixpunktsatz für ultrametrische Räume*, Results in Mathematics **18** (1990), 178–186
- [PR] Priß-Crampe, S. – Ribenboim, P.: *Fixed Points, Combs and Generalized Power Series*, Abh. Math. Sem. Hamburg **63** (1993), 227–244
- [R1] Rosenlicht, M.: *Differential valuations*, Pacific J. Math. **86** (1980), 301–319
- [R2] Rosenlicht, M.: *On the value group of a differential valuation*, Amer. J. Math. **191** (1979), 258–266

The Fields Institute, 222 College Street, Toronto, Ontario M5T 3J1, Canada
email: fkuhlman@fields.utoronto.ca